

Taq PCR Kit

Description: An ultrapure recombinant thermostable *Taq* DNA polymerase obtained by high level expression of the *Taq* DNA polymerase gene in *E. coli*. It is licensed and optimized for use in the Polymerase Chain Reaction (PCR) process.

Enzyme Source: *Thermus aquaticus*

Concentration: 5 units/ μ l

Unit Definition: One unit of enzyme is defined as the amount that will incorporate 10nmoles of dNTPs into acid insoluble material in 30 minutes at 74°C under the analysis conditions below.

Associated Activities: *Taq* DNA polymerase has 5' to 3' polymerization and exonuclease activity but lacks 3' to 5' exonuclease activity (proofreading).

Kit Contents:

Vial (cap color)	Pack Size	
	A	B
<i>Taq</i> DNA Polymerase (clear)	50 μ l	10 x 50 μ l
Buffer IV containing 15mM MgCl ₂ (red)	1.25ml	10 x 1.25ml
dNTP Mix (brown)	0.5ml	3 x 1.6ml

<u><i>Taq</i> DNA</u>	100mM	KCl
<u>Polymerase:</u>	20mM	Tris-HCl, pH 8.0 (at 25°C)
	0.1mM	EDTA (ethylenediaminetetraacetic acid)
	1mM	DTT (dithiothreitol)
	0.5%	Tween® 20
	0.5%	Nonidet® P40
	50% (v/v)	Glycerol
<u>Reaction</u>	750mM	Tris-HCl, pH 8.8 (at 25°C)
<u>Buffer IV</u>	200mM	(NH ₄) ₂ SO ₄
<u>+MgCl₂ (10X):</u>	0.1% (v/v)	Tween® 20

For Research Purposes Only

1

dNTP Mix:	5mM	dATP
	5mM	dCTP
	5mM	dGTP
	5mM	dATP

Storage Conditions:

Store at -20°C until ready for use. *Taq* DNA polymerase is stable for a minimum of 12 months. The reagents can be stored at 4°C for up to 1 month. Avoid repeated freeze thawing. Shipped on ice within the UK and on dry ice for international and within the US.

Example of Protocol:

Mix and spin down the solutions prior to use

	Volume	Final Concentration 1X
<i>Taq</i> DNA Polymerase (5U/μl)	0.125μl	0.625 U
10X Reaction Buffer IV + MgCl ₂ (15mM)	2.5μl	1X with 1.5mM MgCl ₂
dNTP Mix (20mM)	1μl	0.2mM of each nucleotide
Primer forward (10μM each)	1.25μl*	0.5μM*
Primer reverse (10μM each)	1.25μl*	0.5μM*
Water (PCR Grade)	Variable	
DNA Template	0.5 - 10μl	0.5 - 125ng
Total volume	25μl	

*Scale up or down the volume and concentration as appropriate

Example of Program:

	Temp.	Time	Number of cycle
Initial Denaturation	94°C	2 min	1 cycle
Denaturation	94°C	20 sec	30 to 40 cycles
Annealing	50-65°C	30 sec	
Extension**	72°C	60 sec	
Final Extension	72°C	5 min	1 cycle

**Increase length of time in proportion to size of amplicon, *Taq* DNA Polymerase extends at approximately 1000 bp/min.

Analysis Conditions:	25mM	TAPS, pH 9.3 (at 25°C)
	50mM	[tris-(hydroxymethyl)-methyl-amino-propane sulfonic acid, sodium salt]
	2mM	KCl
	1mM	MgCl ₂
	250µM	β-mercaptoethanol
	250µM	of each: dCTP, dGTP, dTTP
	1.25µg/µl	[³ H] dATP (0.05 Ci/mmol)
		activated salmon sperm DNA

Water added to a total volume of 50µl. Incubated at 74°C for 10 minutes.

Ordering Information:	AB-1192/A/N	Taq PCR Kit	250 units
	AB-1192/B/N	Taq PCR Kit	2500 units

All sizes are supplied with 10X Reaction Buffer IV containing 15mM MgCl₂ and a 20mM dNTP Mix.

Troubleshooting

For troubleshooting, see www.abgene.com/troubleshoot.asp or contact Thermo Fisher Scientific (ABgene) TechSupport at abgene.techsupport@thermofisher.com

UK TechSupport, call +44 (0) 1372 840 410

For all other regions, please contact your local Thermo Fisher Scientific (ABgene) office / distributor.

Use of this product is covered by one or more of the following US patents and corresponding patent claims outside the US: 5,079,352, 5,789,224, 5,618,711, 6,127,155 and claims outside the US corresponding to US Patent No. 4,889,818. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. No right under any other patent claim (such as the patented 5' Nuclease Process claims in US Patents Nos. 5,210,015 and 5,487,972), no right to perform any patented method, and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is conveyed expressly, by implication, or by estoppel. This product is for research use only. Diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained by contacting the Director of Licensing, Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404, USA.

For Research Purposes Only

3